

Preliminary communication

9-(β -D-Apio-L-furanosyl)-2-chloroadenine*

F. PERINI**, F. A. CAREY***, and L. LONG, Jr.

Pioneering Research Laboratory, U.S. Army Natick Laboratories, Natick, Massachusetts 01760 (U.S.A.)

(Received June 10th, 1969; in revised form July 10th, 1969)

The preparation of C-3'-modified purine nucleosides has both chemical and pharmacological interest in view of the recent syntheses of such compounds. These systems often exhibit antibiotic and/or antitumor activity. Some of these compounds are cordycepin (3'-deoxyadenosine)^{1,2}, puromycin³, 3'-C-alkylerythrofuranosyl nucleosides^{4,5} and, most recently, a nogalose (a 3'-C-methyl-D-allopyranose) nucleoside⁶. Our program on the chemistry of apiose⁷ includes the production of nucleosides from this compound, which is one of the most widespread among naturally occurring branched-chain sugars.

We now wish to report the first synthesis of an apiosyl nucleoside of unequivocal structure.

The configuration at C-3 and C-5 of the D-apio-L-furanose system was locked by cyclocarbonation. Treatment of 1,2-O-isopropylidene- α -D-apio-L-furanose⁸ (1) with *N,N'*-carbonyldiimidazole in tetrahydrofuran gave (92%) 3,5-O-carbonyl-1,2-O-isopropylidene- α -D-apio-L-furanose† (2), m.p. 113–114°; $[\alpha]_D^{22} + 63.7^\circ$, (*c* 1.9, chloroform); $\lambda_{\text{max}}^{\text{Nujol}} 5.52 \mu\text{m}$ (–O–CO–O–). Sulfuric acid-catalyzed acetolysis of 2 in acetic acid–acetic anhydride yielded (42–59%) crystalline 1,2-di-O-acetyl-3,5-O-carbonyl- α and β -D-apio-L-furanose (3), m.p. 141.5–145°; $[\alpha]_D^{23} + 38.6^\circ$ (*c* 1.1, chloroform); $\lambda_{\text{max}}^{\text{Nujol}} 5.50 \mu\text{m}$ (–O–CO–O–) and 5.71 μm (OAc). The n.m.r. spectrum†† (chloroform-*d*) included: 6.44 (0.3-proton doublet, $J_{1,2}$ 4.5 Hz, H-1, 30% α -anomer) and 6.13 (0.7-proton singlet, H-1, 70% β -anomer). A syrupy product, partially characterized as 1,1,2,4-tetra-O-acetyl-3,5-O-carbonylapiose (4). (~40–49%) $\lambda_{\text{max}} 5.52 \mu\text{m}$ (–O–CO–O–) and 5.71 μm (OAc); n.m.r. (chloroform-*d*): 6.96 (1-proton doublet, $J_{1,2}$ 3 Hz, H-1), was also formed during acetolysis of 2.

Fusion of 2,6-dichloropurine (DCP) and 3 in the presence of dichloroacetic acid (Cl_2CHCOOH)⁹ at 160° was successful. This technique, as found in other systems^{10,11}, gave a mixture (58%, after silica gel column chromatography) of α,β -anomers, 2,6-dichloro-9-(2'-O-acetyl-3', 5'-O-carbonyl- α - and β -D-apio-L-furanosyl)purine (5a and 5b), m.p.


*Paper VI of a series of publications from this laboratory concerning the chemistry of apiose.

**PRL Postdoctoral Research Fellow, 1967–1969.

***Present address: Department of Chemistry, University of Virginia, Charlottesville, Virginia 22901, U.S.A.

†Acceptable elemental analyses were obtained for all crystalline compounds reported. Melting points are uncorrected.

††100 MHz, with chemical shifts reported in δ units from a tetramethylsilane internal standard.



115–120°; n.m.r. (chloroform-*d*): 6.68 (0.32-proton doublet, $J_{1',2'}$ 5.5 Hz, H-1', 32% α -anomer)¹², and 6.26 (0.68-proton doublet, $J_{1',2'}$ 2.5 Hz, H-1', 68% β -anomer)¹². The chromatographic mobilities of 5a and 5b were different but too similar to allow preparative-scale separation.

Condensation of DCP with 3 in nitromethane at reflux^{13, 14}, in the presence of dichloroacetic acid, gave only the pure branched-chain sugar nucleoside 5a (51%, after silica gel column chromatography); m.p. 177–177.5°; $[\alpha]_D^{25} + 55.0^\circ$ (*c* 0.7, chloroform); $\lambda_{\text{max}}^{\text{Nujol}}$ 5.49 (O-CO-O-), 5.70 (OAc) and 6.25, 6.41 μm (C=N, C=C); $\lambda_{\text{max}}^{\text{EtOH}}$ (pH 6) 255 shoulder (ϵ , 5660) and 274.5 nm (ϵ , 9650). Formation of 5a in nitromethane, a solvent which promotes β -D anomer production in glycoside synthesis¹⁵, was virtually stereospecific, as shown by the n.m.r. data (chloroform-*d*): 8.40 (1-proton singlet, H-8), 6.20 (1-proton doublet, $J_{1',2'}$ 2.5 Hz, H-1'), 5.67 (1-proton doublet, $J_{1',2'}$ 2.5 Hz, H-2'), 4.70, 4.18 (2-proton AB quartet, $J_{4',4'}$ 11 Hz, H-4'); 4.69, 4.40 (2-proton AB quartet, $J_{5',5'}$ 10 Hz, H-5'), and 2.22 (3-proton singlet, OAc). The β -D anomeric configuration was assigned to 5a due to the observation of a narrow spacing (2.5 Hz) of the doublet¹² ascribable to H-1'. Predominant formation of a *trans* nucleoside would be expected from 2-O-acyl group participation¹⁶.

Other Lewis acids, such as sulfamic acid¹⁴ (76%) and boron trifluoride etherate (57%), also catalyzed the highly stereoselective formation of 5a. Heating DCP with 3 in nitromethane for 1.5 day at 120° without catalyst, however, gave no product. It is known¹⁷ that, in acidic media, DCP is hydrolyzed to xanthine. Under dry conditions,

during the synthesis of **5a** in the presence of dichloroacetic acid 27% of the DCP was converted into xanthine.

The convenient synthesis of **5a** opens the way to the production of various apiose nucleosides modified on the purine ring¹⁸. Ammonolysis of **5a** with saturated methanolic ammonia in a sealed tube^{18, 19}, followed by picrate formation and liberation of the base with an aqueous suspension of AG 1-X4 (CO₃²⁻) anion exchange resin gave the useful deblocked apioside (49%), 9-(β -D-apio-L-furanosyl)-2-chloroadenine (**6**), m.p. 137–138°; $[\alpha]_D^{26}$ -16.0° (c 0.5, methanol); $\lambda_{\text{Nujol}}^{\text{max}}$ 3.00, 3.20 (OH, NH), and 6.02, 6.27, 6.30 μm (NH, C=N, C=C); $\lambda_{\text{max}}^{\text{EtOH}}$ (pH 6) 266 nm (ϵ , 15300); n.m.r. (deuterium oxide): 8.60 (1-proton singlet, H-8), 6.34 (1-proton doublet, $J_{1', 2'}$ 2 Hz, H-1')¹², 4.90 (1-proton doublet, $J_{1', 2'}$ 2 Hz, H-2'), 4.75 (2-proton singlet, H-5'), and 4.67, 4.63 (2-proton singlets, H-4').

Work on the synthesis of several derivatives of **5a** and **6** is in progress, and will be reported in full elsewhere.

ACKNOWLEDGMENT

The authors thank Mr. F. H. Bissett and Dr. R. C. Chalk for the measurement of n.m.r. spectra.

REFERENCES

- 1 E. Walton, F. W. Holly, G. E. Boxer, R. F. Nutt, and S. R. Jenkins, *J. Med. Chem.*, **8** (1965) 659.
- 2 J. Prokop and D. H. Murray, *J. Pharm. Sci.*, **54** (1965) 1468.
- 3 J. J. Fox, K. A. Watanabe, and A. Bloch, *Progr. Nucleic Acid Res. and Mol. Biol.*, **5** (1966) 253.
- 4 E. J. Reist, D. F. Calkins, and L. Goodman, *J. Amer. Chem. Soc.*, **90** (1968) 3852.
- 5 R. F. Nutt, M. J. Dickinson, F. W. Holly, and E. Walton, *J. Org. Chem.*, **33** (1968) 1789.
- 6 G. B. Howarth, W. A. Szarek, and J. K. N. Jones, *Can. J. Chem.*, **46** (1968) 3691.
- 7 D. H. Ball, F. A. Carey, I. L. Klundt, and L. Long, Jr., *Carbohydr. Res.*, **10** (1969) 121 and references cited therein.
- 8 F. A. Carey, D. H. Ball, and L. Long, Jr., *Carbohydr. Res.*, **3** (1966) 205.
- 9 Y. H. Pan, R. K. Robins, and L. B. Townsend, *J. Heterocycl. Chem.*, **4** (1967) 246.
- 10 W. W. Lee, A. P. Martinez, G. L. Tong, and L. Goodman, *Chem. Ind.*, (London), (1963) 2007.
- 11 J. A. M. Montgomery and K. Hewson, *Chem. Commun.*, (1969) 15.
- 12 J. A. Montgomery and K. Hewson, *J. Med. Chem.*, **11** (1968) 48, and references cited therein.
- 13 M. Honjo, S. Furukawa, and K. Kobayashi, Japanese Patent 42-11077 (June 20, 1967); *Chem. Abstr.*, **67** (1967) 188958r.
- 14 Y. Ishido, H. Tanaka, T. Yoshino, M. Sekiya, K. Iwabuchi, and T. Sato, *Tetrahedron Lett.*, (1967) 5245.
- 15 P. F. Lloyd and G. P. Roberts, *J. Chem. Soc.*, (1963) 2962.
- 16 T. Sato, T. Simadate, and Y. Ishido, *Nippon Kagaku Zasshi*, **81** (1960) 1440.
- 17 E. H. Rodd, ed., *Chemistry of Carbon Compounds, Heterocyclic Compounds*, Vol. IVc, Elsevier, Amsterdam, 1960, p. 1672.
- 18 H. J. Schaeffer and H. J. Thomas, *J. Amer. Chem. Soc.*, **80** (1958) 3738.
- 19 Y. Ishido, Y. Kikuchi, and T. Sato, *Nippon Kagaku Zasshi*, **86** (1965) 240; (also, T. Sato, *Synth. Procedures Nucleic Acid Chem.*, **1** (1968) 267)

Carbohydr. Res., **11** (1969) 159–161